

Subvenite[®]

(lamotrigine tablets, USP)

Simplify Treatment
from the Start

STEP 1

Print, or download on your smart phone, this Subvenite[®] Patient Savings Card.

STEP 2

Present your printed, or downloaded, card to the pharmacist for immediate savings.

If your insurance covers Subvenite[®] Starter Kits, most patients pay \$5 for any Subvenite[®] Starter Kit.



Inform your pharmacist of your preference for the Subvenite[®] brand.



Check the label on your prescription to be sure it says Subvenite[®].



When filling a prescription for Subvenite[®], make sure your kit look like those shown. If not, call your doctor or talk to your pharmacist.



The Pharmaceutical Social Enterprise[®]

400 E Diehl Rd, Suite 400 • Naperville, IL 60563 | info@owppharma.com | 1-800-273-6729 | owppharma.com
©2020 OWP Pharmaceuticals, Inc. All rights reserved. Printed in U.S.A. OIWOS1050V4/6/20

Patient Instructions: You must present this card to the pharmacist along with your prescription to participate in this program. If you have any questions regarding your eligibility, or if you wish to discontinue your participation, call the Subvenite[®] co-pay assistance program at 1-800-273-6729 (9:00 AM - 5:00 PM CST, Monday - Friday). When you use this card, you are certifying that you understand and agree to comply with the program restrictions, regulations, and terms and conditions.

Pharmacist Instructions: When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under any federal, state, or other governmental programs for this prescription and will otherwise comply with the terms of this offer.

- Transmit claim online to RxBIN# 018844
- Commercially Insured Patients: Submit claim to patient's primary insurance first, then submit this offer as a secondary claim, as a Coordination of Benefits using other coverage code type 08.
- Cash-Paying Patients: Submit as primary claim using other coverage code 0 or 01.
- For assistance filing this claim, call Help Desk at 1-855-282-4888.

Program Restrictions: Not eligible if prescriptions are paid by any state or federally funded programs, including, but not limited to Medicare, Medicaid, VA, DOD, Tricare or Medigap. Void where prohibited by law. Not valid for a Subvenite[®] prescription reimbursed in full by any third-party payer. May not be combined with any other coupon, discount, savings card or other offer. May not be accepted at all pharmacies. Starter Kit: Limited to one prescription per person for any 35-day period. The maximum benefit is \$50 per 35-day starter kit prescription. No substitutions permitted. Maintenance Dosage: Limited to one prescription per person for any 30-day period. The maximum benefit is \$5 per 30-day prescription.

This offer may be terminated, rescinded, revoked or amended by OWP Pharmaceuticals, Inc. at any time, without notice.



The Pharmaceutical Social Enterprise[®]

400 E Diehl Rd, Suite 400 • Naperville, IL 60563 | info@owppharma.com | 1-800-273-6729 | owppharma.com

Subvenite[®]
(lamotrigine tablets, USP)

Patient Savings Card

RxBIN: 018844
RxPCN: 3F
RxGRP: FCSUBCP17
RxID: SUBW7281798

Eligible commercial patients are responsible for \$20 or less out-of-pocket expense per Starter Kit and per 30-day maintenance supply.

Cash-paying patients are estimated to pay approximately \$50 for a Starter Kit and approximately \$30 out-of-pocket expense per 30-day maintenance supply.

Subvenite[®] is a registered trademark of OWP Pharmaceuticals, Inc.
©2020 OWP Pharmaceuticals, Inc. Naperville, IL 60563
All rights reserved. Printed in U.S.A.



OUT CARD OUT AND FOLD IN HALF

MEDICATION GUIDE
SUBVENITE (Sub-VE-nite) lamotrigine tablets, USP

What is the most important information I should know about SUBVENITE?

1. SUBVENITE may cause a serious skin rash that may cause you to be hospitalized or even cause death.
There is no way to tell if a mild rash will become more serious. A serious skin rash can happen at any time during your treatment with SUBVENITE, but is more likely to happen within the first 2 to 8 weeks of treatment. Children and teenagers aged between 2 and 17 years have a higher chance of getting this serious skin rash while taking SUBVENITE.
The risk of getting a serious skin rash is higher if you:

- take SUBVENITE while taking valproate [DEPAKENE (valproic acid) or DEPAKOTE (divalproex sodium)].
- take a higher starting dose of SUBVENITE than your healthcare provider prescribed.
- increase your dose of SUBVENITE faster than prescribed.

Call your healthcare provider right away if you have any of the following:

- a skin rash
- blistering or peeling of your skin
- hives
- painful sores in your mouth or around your eyes

These symptoms may be the first signs of a serious skin reaction. A healthcare provider should examine you to decide if you should continue taking SUBVENITE.

2. Other serious reactions, including serious blood problems or liver problems.
SUBVENITE can also cause other types of allergic reactions or serious problems that may affect organs and other parts of your body like your liver or blood cells. You may or may not have a rash with these types of reactions. Call your healthcare provider right away if you have any of these symptoms:

- fever
- frequent infections
- severe muscle pain
- swelling of your face, eyes, lips, or tongue
- swollen lymph glands
- unusual bruising or bleeding
- weakness, fatigue
- yellowing of your skin or the white part of your eyes

3. Like other antiepileptic drugs, SUBVENITE may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.
Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempt to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Do not stop SUBVENITE without first talking to a healthcare provider.

- Stopping lamotrigine suddenly can cause serious problems.
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions in myself or a family member?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

- Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

4. SUBVENITE may cause aseptic meningitis, a serious inflammation of the protective membrane that covers the brain and spinal cord.
Call your healthcare provider right away if you have any of the following symptoms:

- headache
- fever
- nausea
- vomiting
- stiff neck
- rash
- unusual sensitivity to light
- muscle pains
- chills
- confusion
- drowsiness

Meningitis has many causes other than SUBVENITE, which your doctor would check for if you developed meningitis while taking SUBVENITE.

SUBVENITE can cause other serious side effects. For more information ask your healthcare provider or pharmacist. Tell your healthcare provider if you have any side effect that bothers you. Be sure to read the section below entitled "What are the possible side effects of SUBVENITE?"

5. People prescribed SUBVENITE have sometimes been given the wrong medicine because many medicines have names similar to SUBVENITE, so always check that you receive SUBVENITE.
Taking the wrong medication can cause serious health problems. When your healthcare provider gives you a prescription for SUBVENITE:

- Make sure you can read it clearly.
- Talk to your pharmacist to check that you are given the correct medicine.
- Each time you fill your prescription, check the tablets you receive against the pictures of the tablets below.

These pictures show the distinct wording, colors, and shapes of the tablets that help to identify the right strength of SUBVENITE tablets. Immediately call your pharmacist if you receive a SUBVENITE tablet that does not look like one of the tablets shown below, as you may have received the wrong medication.

SUBVENITE (lamotrigine) tablets, USP

Tablet Strength	SUBVENITE Dimensional Drawing
25 mg	
100 mg	
150 mg	
200 mg	

What is SUBVENITE?

- SUBVENITE is a prescription medicine used:
 - together with other medicines to treat certain types of seizures (partial-onset seizures, primary generalized tonic-clonic seizures, generalized seizures of Lennox-Gastaut syndrome) in people aged 2 years and older.
 - alone when changing from 1 other medicine used to treat partial-onset seizures in people aged 16 years and older.
 - for the long-term treatment of bipolar I disorder to lengthen the time between mood episodes in people who have been treated for mood episodes with other medicine.
- It is not known if SUBVENITE is safe or effective in people younger than 18 years with mood episodes such as bipolar disorder or depression.
- It is not known if SUBVENITE is safe or effective when used alone as the first treatment of seizures.

- It is not known if SUBVENITE is safe or effective for people with mood episodes who have not already been treated with other medicines.
- SUBVENITE should not be used for acute treatment of manic or mixed mood episodes.

Do not take SUBVENITE:
if you have had an allergic reaction to lamotrigine or to any of the inactive ingredients in SUBVENITE. See the end of this leaflet for a complete list of ingredients in SUBVENITE.

Before taking SUBVENITE, tell your healthcare provider about all of your medical conditions, including if you:

- have had a rash or allergic reaction to another antiseizure medicine.
- have or have had depression, mood problems, or suicidal thoughts or behavior.
- have had aseptic meningitis after taking SUBVENITE.
- are taking oral contraceptives (birth control pills) or other female hormonal medicines. Do not start or stop taking birth control pills or other female hormonal medicine until you have talked with your healthcare provider. Tell your healthcare provider if you have any changes in your menstrual pattern such as breakthrough bleeding. Stopping these medicines while you are taking SUBVENITE may cause side effects (such as dizziness, lack of coordination, or double vision). Starting these medicines may lessen how well SUBVENITE works.
- are pregnant or plan to become pregnant. It is not known if SUBVENITE will harm your unborn baby. If you become pregnant while taking SUBVENITE, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of antiepileptic drugs during pregnancy.
- are breastfeeding. SUBVENITE passes into breast milk and may cause side effects in a breastfed baby. If you breastfeed while taking SUBVENITE, watch your baby closely for trouble breathing, episodes of temporarily stopping breathing, sleepiness, or poor sucking. Call your baby's healthcare provider right away if you see any of these problems. Talk to your healthcare provider about the best way to feed your baby if you take SUBVENITE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. SUBVENITE and certain other medicines may interact with each other. This may cause serious side effects.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take SUBVENITE?

- Take SUBVENITE exactly as prescribed.
- Your healthcare provider may change your dose. Do not change your dose without talking to your healthcare provider.
- Do not stop taking SUBVENITE without talking to your healthcare provider. Stopping SUBVENITE suddenly may cause serious problems. For example, if you have epilepsy and you stop taking SUBVENITE suddenly, you may have seizures that do not stop. Talk with your healthcare provider about how to stop SUBVENITE slowly.
- If you miss a dose of SUBVENITE, take it as soon as you remember. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. **Do not take 2 doses at the same time.**
- If you take too much SUBVENITE, call your healthcare provider or your local Poison Control Center or go to the nearest hospital emergency room right away.
- You may not feel the full effect of SUBVENITE for several weeks.
- If you have epilepsy, tell your healthcare provider if your seizures get worse or if you have any new types of seizures.
- Swallow SUBVENITE whole.
- If you have trouble swallowing SUBVENITE

tablets, tell your healthcare provider because there may be another form of lamotrigine you can take.

- If you receive SUBVENITE in a blister-pack, examine the blister-pack before use. Do not use if blisters are torn, broken, or missing.

What should I avoid while taking lamotrigine?
Do not drive, operate machinery, or do other dangerous activities until you know how lamotrigine affects you.

What are the possible side effects of SUBVENITE?
SUBVENITE can cause serious side effects. See "What is the most important information I should know about SUBVENITE?"
Common side effects of SUBVENITE include:

- dizziness
- tremor
- headache
- sleepiness
- back pain
- nausea, vomiting
- diarrhea
- tiredness
- insomnia
- dry mouth
- stuffy nose
- sore throat
- seasonal flu

These are not all the possible side effects of SUBVENITE. For more information, ask your healthcare provider or pharmacist.
Call your doctor for medical advice about side effects. You may report side effects to FDA 1-800-FDA-1088.

How should I store SUBVENITE?

- Store SUBVENITE at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].
- Keep SUBVENITE and all medicines out of the reach of children.

General information about the safe and effective use of SUBVENITE.
Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use SUBVENITE for a condition for which it was not prescribed. Do not give SUBVENITE to other people, even if they have the same symptoms that you have. It may harm them.
If you take a urine drug screening test, SUBVENITE may make the test result positive for another drug. If you require a urine drug screening test, tell the healthcare professional administering the test that you are taking SUBVENITE.
You can ask your healthcare provider or pharmacist for information about SUBVENITE that is written for healthcare professionals.
For more information, call 1-800-273-6729.

What are the ingredients in SUBVENITE? SUBVENITE (lamotrigine) tablets, USP
Active ingredient: lamotrigine, USP.
Inactive ingredients: lactose monohydrate; magnesium stearate; microcrystalline cellulose; povidone; and sodium starch glycolate.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

torrent PHARMA
Manufactured by:
TORRENT PHARMACEUTICALS LTD., INDIA.
Manufactured for:
OWPPHarmaceuticals, Inc., 400E. Dieh Road, Suite 400, Naperville, IL 60563.
OWOSSUBMG1019 Revised October 2019 8076981

